

IS THE GLASSY-WINGED SHARPSHOOTER PARASITOID *GONATOCERUS MORRILLI* ONE SPECIES OR A COMPLEX OF CLOSELY RELATED SIBLING SPECIES?

Project Leaders:

Mark Hoddle
Dept. of Entomology
University of California
Riverside, CA 92521

Richard Stouthamer
Dept. of Entomology
University of California
Riverside, CA 92521

Cooperator:

Serguei Triapitsyn
Dept. of Entomology
University of California
Riverside, CA 92521

Reporting period: The results reported here are from work conducted from July 2004 to October 2004.

INTRODUCTION

This is a new proposal that was officially funded in July 2004. This project objective is to determine the status of different *Gonatocerus morrilli* populations. We intend to use three approaches to determine the species identity of different *G. morrilli* populations: (1) Reassessment of key morphological features using scanning electron microscopy to determine if subtle morphological differences exist between *G. morrilli* populations which could possibly indicate species differences (Triapitsyn to conduct this work). (2) Conduct mating compatibility studies to determine if different populations of *G. morrilli* are reproductively isolated, or if mating occurs, whether offspring are viable thereby defining species groups on the basis of successful interbreeding (Hoddle). (3) To determine if molecular differences exist between *G. morrilli* populations collected from different regions by comparing mitochondrial and ribosomal DNA sequences. Molecular dissimilarities of key regions could potentially indicate the existence of different species (Stouthamer). Results from these three areas (morphology, behavior, and molecular) of investigation will be evaluated together to determine whether *G. morrilli* as it is currently viewed is a valid species or whether it is an aggregate of morphologically similar cryptic species.

A classical biological control program is currently underway for glassy-winged sharpshooter (GWSS), which is an exotic pest in California. The native range of GWSS is the southeastern United States and northeastern Mexico (Triapitsyn & Phillips, 2000). GWSS is thought to have invaded California around 1990 as egg masses that were accidentally imported on ornamental plants from Florida. Species of GWSS egg parasitoids not present in California are currently being prospected for in the native range of GWSS. Promising candidate natural enemy species that attack eggs are being imported and released in California for GWSS control (Triapitsyn et al., 1998; Triapitsyn & Hoddle, 2001). Interestingly, one species of egg parasitoid associated naturally with GWSS in California, *Gonatocerus morrilli* (Howard) (Hymenoptera: Mymaridae), is also widely distributed in the home range of GWSS, but at the time of its initial discovery in California, *G. morrilli* had not been intentionally released here and was thought to be native to California. A potential host for *G. morrilli* in California prior to the arrival of GWSS could have been the native *Homalodisca liturata* (Ball) which has had unidentified *Gonatocerus* spp. reared from its egg masses collected in the San Diego area (Powers, 1973). The presence of *G. morrilli* in Riverside in 1980-1984 has been documented (Huber 1988). *Gonatocerus morrilli* is now the second most important natural enemy of GWSS egg masses in California (Al-Wahaibi, 2004).

The success and failure of a number of biological control projects against insect pests and weeds has hinged on the correct taxonomic identification of the target and its natural enemies (Gordh and Beardsley, 1999). Incorrect understanding of the taxonomy and subsequent interrelationships between the target and its natural enemy guild are serious impediments to an efficacious biological control program. For example, *Trichogramma minutum* and *T. platneri* are important commercially available biological control agents that are morphologically indistinguishable but reproductively incompatible (Nagarkatti, 1975). Experimental work and subsequent modeling with these two species of *Trichogramma* has indicated that because pre-mating isolation mechanisms are absent (e.g., pre-mating courtship behaviors that prevent coupling of males and females from different species) severe negative effects on biological control can occur. Negative effects manifest themselves because females that mate with males from different species fail to produce female offspring. This occurs because *Trichogramma* like *Gonatocerus* are haploid-diploid parasitic Hymenoptera. In this haplo-diploid system, fertilized eggs produce female offspring and unfertilized eggs produce male offspring. In situations where incompatible interspecies matings are occurring both species fail to produce females and the potential population growth of both parasitoid species is reduced to levels below the growth rate expected for either species in the absence of the other (Stouthamer et al., 2000).

If different populations of morphologically similar *G. morrilli* from Florida, Louisiana, Texas, and Mexico are indeed valid species that lack pre-mating isolation mechanisms, then the current biological control program against GWSS in California that is attempting to establish these new agents may reduce the current level of control achieved by the precinctive populations of *G. morrilli* in California. This could occur because of male-biased offspring production resulting from incompatible matings across species. The rationale for introducing new strains or races of *G. morrilli* into California is based on the idea that different biotypes of this parasitoid may exist and fill niches not currently occupied by the strain of *G. morrilli* already present in California.

In this grant we propose to determine if geographically distinct populations of *G. morrilli* are part of one continuous interbreeding population or if populations of *G. morrilli* are separate species that can't be easily separated on the basis of

currently employed morphological characters. To do this we intend to combine three separate approaches to determine the species identity of different *G. morrilli* populations: First, we'll reassess key morphological features used to characterize *G. morrilli* with scanning electron microscopy to determine if subtle morphological differences exist between *G. morrilli* populations which could possibly indicate species differences. Such differences - should they exist - may not be easily observed with light microscopy. Second, we'll conduct mating compatibility studies to determine if different populations of *G. morrilli* are reproductively isolated, or if mating occurs, whether offspring are viable thereby defining species groups on the basis of successful interbreeding. Third, we'll determine if molecular differences exist between different *G. morrilli* populations by comparing mitochondrial and ribosomal DNA sequences. Molecular dissimilarities of key regions could potentially indicate the existence of different species, and at the same time allow their identification. Results from these three areas (morphology, behavior, and molecular avenues) of investigation will be evaluated together to determine whether *G. morrilli* as it is currently viewed is a valid species or whether it is an aggregate of morphologically indistinguishable cryptic species.

RESULTS

This project has not commenced. The reason for this is that the recruitment of the post-doc has taken some time. We expect the post-doc to be on-line in early December 2004. We will be formally requesting a no-cost extension for this project.

FUNDING AGENCIES

Funding for this project was provided by the University of California Pierce's Disease Grant Program.