

# DEVELOPMENT OF SSR MARKERS FOR GENOTYPING AND ASSESSING THE GENETIC DIVERSITY OF *XYLELLA FASTIDIOSA* IN CALIFORNIA

## Project Leaders:

Hong Lin  
Crop Diseases, Pests & Genetics  
SJVAC, USDA-ARS  
Parlier, CA 93648

Andrew Walker  
Dept. of Viticulture & Enology  
University of California  
Davis, CA 95616

## Collaborator:

Edwin Civerolo  
SJVAC, USDA-ARS  
Parlier, CA 93648

**Reporting period:** The results reported here are from work conducted from March 2004 to September 2004.

## ABSTRACT

Recently available genomic sequences of four *Xylella fastidiosa* strains (PD, CVCD, ALSD and OLS) facilitate genome wide searches for identifying Simple Sequence Repeat (SSR) loci. Sixty SSR loci were selected for SSR marker development. We designed and validated 34 SSR primers with good reliability and specificity. These SSR primers showed various levels of polymorphism with average 11.3 alleles per locus among 43 *Xylella fastidiosa* isolates. These multi-locus SSR markers, distributed across the entire genome, are a useful tool for pathogen genotyping, population genetics and molecular epidemiology studies.

## INTRODUCTION

*Xylella fastidiosa* (*Xf*) causes economically important diseases that results in significant losses in several agricultural, horticultural and landscape crops, including grape Pierce's disease (PD), almond leaf scorch disease (ALSD), citrus variegated chlorosis disease (CVCD) and oleander leaf scorch disease (OLS). Recent introduction and establishment of the invasive and more effective vector, the Glassy-winged Sharpshooter (*Homalodisca coagulata*, GWSS) has had a great impact on the California grape industry. Host plant resistance is a critical component of integrated crop management. If this insect becomes widely established, the use of resistant varieties may become the most reliable and effective way to control PD. However, the durability of resistant grape plants depends upon the variability and adaptability of the pathogen population and its interaction with the resistance genes of plants. Most resistance studies are performed by screening against a subpopulation of a given pathogen, and neglect that fact that changes in pathogen population structure that may lead to resistance breakdown.

It is clear that pathogen populations with a high evolutionary potential are more likely to overcome host genetic resistance than pathogen populations with a low evolutionary potential (MacDonald and Linde, 2002). The risk becomes even greater with the recent establishment of a more effective vector, the GWSS, which dramatically increases the dispersal of *Xf* genes/genotypes. In California, information regarding the population structure and genetic diversity, as well as the genetic evolutionary and epidemiological relationships, among *Xf* strains in agricultural populations is not clear. In order to develop effective management strategies, it is critical to understand pathogen population structure and genetic diversity in the agricultural ecosystem. A tool is needed that is capable of precisely, powerfully, easily analyzing *Xf* diversity and genotyping strains. We developed multi-locus DNA markers to fill this need.

## OBJECTIVES

1. Perform genome-wide sequence analysis to identify Simple Sequence Repeat (SSR) loci from four *Xf* genomic sequencing databases (PD, CVCD, ALSD and OLS). Design and develop multi-locus SSR markers.
2. Analyze genetic diversity and population structures of PD *Xf* statewide. Compile a large *Xf* allele frequency database for strain identification.
3. Use the SSR Marker system to examine interactions between hosts and *Xf* including adaptation, host selection and pathogenicity of *Xf* strains

## RESULTS


### *SSR Locus Identification and Primer Design*

1. A genome wide search was performed to identify SSR loci among all four *Xf* strains (CVC 9a5c 2.68Mbp, PD Temecula 2.52Mbp, ALS Dixon 2.67Mbp, and OLS Ann-1 2.63Mbp). Figure 1 shows the distributions of SSR loci among four strains of *Xf*.
2. We used the following criteria to select SSR loci for primer design; a) each locus has single hit per genome and b) each selected locus contains at least 5 or more of repeat unit lengths.


3. Sequence alignment was then performed to remove redundant loci and to identify conserved flanking sequence regions across four strains for priming sites between 100-200 bp up/down stream of each repeat locus. This step ensures that primers designed will work for all *Xf* strains.
4. BLAST analysis was performed to examine each selected locus against more than 300 microbial genomes in GeneBank to ensure selected loci are unique. No significant hits were found (E value  $< e^{-30}$ ).
5. All SSR primers were designed using the same parameters (50% GC,  $T_m = 60^\circ\text{C}$ , primer length  $\approx 20\text{bp}$ , and self dimer/cross dimer  $\Delta G = -5 \text{ kcal/mol}$ ). This facilitated SSR primer validation and should facilitate scaling up to multiplex PCR formats in future.
6. Based on the criteria and conditions above, 50 primers passed the *in silico* validation test.
7. We further evaluated 50 SSR primers using 43 *Xf* isolates collected from grape, citrus, almond and oleander hosts (see Table 1). In this study, we used thirty-four primers. The results of 34 SSR markers analyses are illustrated in Figures 2 and 3.

## CONCLUSION

Repetitive DNA is ubiquitous in microbial genomes. It has been shown to be a useful tool for genetic study in prokaryotes (Belkum, et al 1998). Data from our preliminary study demonstrates that this technique works well for discriminating *Xf* strains. This project will provide an accurate and reliable marker system for genotyping, quarantine purposes, genetic diversity analyses, epidemiological analyses and risk assessment studies.




**Figure 1.** Summaries of SSR loci distributions in each strain of *Xylella fastidiosa*. No mono- and di-repeats occur among these four strains. The above illustrates perfect and imperfect simple repeats with repeat unit length = or  $> 5$ .



**Figure 2.** Examples of SSR markers with primers CSSR6 (above) and OSSR9 (below) among 43 *Xylella fastidiosa* isolates separated by 5% of polyacrylamide gel. A, T, C and G are molecular size markers.

Strain Name	Host of Origin	County or state from which strain was collected
PD-1	Grape	Kern, CA
PD-2	Grape	Kern, CA
PD-3	Grape	Kern, CA
PD-4	Grape	Kern, CA
PD-5	Grape	Temecula, CA
PD-6	Grape	Temecula, CA
PD-7	Grape	Temecula, CA
PD-8	Grape	Kern, CA
PD-9	Grape	Kern, CA
PD-10	Grape	Kern, CA
PD-11	Grape	Kern, CA
PD-12	Grape	Baja, CA
PD-13	Grape	Kern, CA
PD-14	Grape	Kern, CA
PD-15	Grape	Napa, CA
PD-16	Grape	Napa, CA
PD-17	Grape	Napa, CA
PD-18	Grape	Napa, CA
PD-19	Grape	Napa, CA
PD-20	Grape	Napa, CA
PD-21	Grape	Napa, CA
PD-22	<b>Grape (Temecula)*</b>	Temecula, CA
CVC-1	Citrus	São Paulo, Brazil
CVC-2	Citrus	São Paulo, Brazil
CVC-3	Citrus	São Paulo, Brazil
CVC-4	Citrus	São Paulo, Brazil
CVC-5	Citrus	São Paulo, Brazil
CVC-6	Citrus	São Paulo, Brazil
CVC-7	Citrus	São Paulo, Brazil
CVC-8	Citrus	São Paulo, Brazil
CVC-9	Citrus	São Paulo, Brazil
CVC-10	<b>Citrus (9a5c)*</b>	Brazil
ALS-1	Almond	Tulare, CA
ALS-2	Almond	Contra Costa, CA
ALS-3	Almond	San Joaquin, CA
ALS-4	Almond	San Joaquin, CA
ALS-5	Almond	San Joaquin, CA
ALS-6	<b>Almond (Dixon)*</b>	Solano, CA
OLS-1	Oleander	Riverside, CA
OLS-2	Oleander	CA
OLS-3	Oleander	CA
OLS-4	<b>Oleander (Ann-1)*</b>	Riverside, CA
OLS-5	Oleander	CA

**Table 1.** 43 *X.fastidiosa* isolates were used for this study. \*Labels in bold are the strains used for genome sequence.



**Figure 3.** Dendrogram shows genetic distance among the 43 isolates in table 1. Data was compiled from 356 alleles generated by 34 SSR loci.

## **REFERENCES**

- Belkum, A., Scherer, S., Alphen, L., and Verbrugh, H. 1998 Short-sequence DNA repeats in prokaryotic genomes. Microbiology and Molecular Biology Review. V. 62, 2:275-293.
- McDonald, B., Linde, C., 2002 Pathogen population genetics, evolutionary potential, and durable Resistance. Annu. Rev. Phytopathol. 40:349-379

## **FUNDING AGENCIES**

Funding for this project was provided by the CDFA Pierce's Disease and Glassy-winged Sharpshooter Board.