

EVALUATION OF BLUE-GREEN SHARPSHOOTER FLIGHT HEIGHT

Project Leader:

Ed Weber
UC Cooperative Extension
Napa, CA 94559

Reporting Period: The results reported here are from work conducted February 2004 to September 2005.

ABSTRACT

Flight heights of blue-green sharpshooters (BGSS) were monitored in the Napa Valley from March through September 2004 and 2005 using pole towers to position yellow sticky cards at heights up to 24 feet. Towers were located adjacent to vineyards at the edge of a riparian zone. Eleven towers were monitored in 2004 and twelve in 2005. Overall trap catches in 2004 were considerably greater than in 2005. For the March-May period, 99% of the catches were made at 15 feet or lower in 2005. For this period in 2004, 76% of the catches were made at 15 feet or lower. This figure increases to 88% with the exclusion of data from one tower in early March (discussed below). These data support the possible use of screen or natural barriers to reduce the number of BGSS entering vineyards in the spring.

INTRODUCTION

Where the BGSS, *Graphocephala atropunctata*, is the primary vector of Pierce's disease (PD), control measures should be aimed at reducing the number of BGSS entering vineyards (4), especially early in the growing season. Early-season infections (March-May) are responsible for most chronic cases of PD (6, 9). Infections resulting from BGSS feeding later in the growing season are not likely to result in PD because most will be eliminated with normal pruning. This is unlike the situation with PD caused by glassy-winged sharpshooter (GWSS) feeding, where chronic infections may occur nearly year-round (1).

Vector control measures in the North Coast include the use of insecticides (4) as well as management of riparian plant communities to reduce the number of favorable BGSS breeding host plants (5).

Another method of reducing vector numbers is to block their flight into vineyards through the use of physical barriers. This could include the use of tall fences made with insect screening materials, as well as natural barriers created by planting dense stands of conifers or other non-host tree species. Both of these approaches are already being employed in a few vineyards in the North Coast, although there are currently no data to show their impacts. The use of barriers has also been suggested as a management tactic to keep GWSS out of vineyards (2).

For barriers to be effective, they would need to block the majority of BGSS from entering vineyards, since small numbers of insects can still lead to significant disease development (8). Unfortunately, little is known about the overwintering behavior of BGSS and its preferred winter plant hosts (7). Therefore, it is not clear how tall a barrier would need to be in order to be effective. Most trapping by both researchers and growers has been done from the ground at the 5-6 foot level.

This project addresses the question of BGSS flight height by installing and monitoring pole towers that can accommodate yellow sticky card trapping up to a height of approximately 24 feet.

OBJECTIVES

1. Evaluate the predominant flight height of BGSS entering vineyards from adjacent riparian habitats through the use of yellow sticky cards positioned at heights from 5 to 24 feet.

RESULTS

Eleven pole towers were installed and monitored in the Napa Valley in 2004, and twelve towers were monitored in 2005. Two of the towers monitored in 2004 were not used in 2005 due to the low number of BGSS trapped at those locations. Three additional towers were installed in 2005. Tower locations covered a distance of approximately 25 miles from the Carneros region in southern Napa County to outskirts of Calistoga at the north end of Napa Valley. Towers were positioned along riparian zones adjacent to vineyards that had a history of PD.

A diagram of a pole tower is shown in Figure 1. Towers were 25 feet in height, constructed from Schedule 40 PVC pipe with a pulley at the top and a rope running through it. Yellow sticky cards were attached to clips on the rope at the following heights: 24 feet, 20 feet, 15 feet and 10 feet. An additional trap at 5 feet was clipped to a metal stake mounted in the ground.




Figure 1. Pole tower diagram.

Towers were installed prior to March 9 in both years. Traps were monitored on a weekly basis through September and numbers of BGSS were recorded. Traps were replaced every two weeks or as needed.

Figure 2 shows the percentage of BGSS trapped at various heights during the early season period of March-May. This is the critical time period in which most infections leading to chronic cases of PD are likely to occur (6, 9).

From March to May 2005, 99% of BGSS were caught at 15 feet and lower. During this same time period in 2004, 76% of BGSS were caught in traps 15 feet and lower. With the exclusion of unusual trap catch data from Tower 10 prior to budbreak in March 2004, this figure rises to 88%. As noted in last year's report (10), Tower 10

was installed adjacent to a Coast Live Oak tree (*Quercus agrifolia*), an evergreen species that was apparently a preferred host plant prior to budbreak of nearby deciduous species. A record heat wave in early March 2004 (70-85°F) led to significant BGSS flight activity in the vicinity of this tree as evidenced by larger numbers of BGSS caught in the upper traps. This was the only case of greater numbers of BGSS in the upper traps compared to the lower traps during the two years of this study at all towers.

Figure 3 shows the percentage of BGSS trapped at various heights during the entire trapping period of March-September for 2004 and 2005. The data in Figures 2 and 3 show similar trends with most BGSS being caught in traps at 15 feet and lower. In 2004, 83% of BGSS were trapped at 15 feet or below. In 2005, 95% were trapped at 15 feet or below. This trend of most BGSS being caught in the lower traps was stronger in 2005 than in 2004, primarily due to the results from Tower 10 noted above.




Figure 2.




Figure 3.

Total BGSS trap catches were over three times greater in 2004 than in 2005. Figures 4 and 5 show the total trap catches for the March-May and March-September periods respectively. The data included in these figures are from the nine towers that were monitored at the same locations in both years. The average cumulative trap catch per tower in 2004 was 36.1 BGSS; in 2005 it was 10.1 BGSS.

The results from this project suggest that a 15-18 foot high barrier could be effective at greatly reducing the number of BGSS entering vineyards. However, previous work with insecticides showed that even with 70-90% reductions in BGSS trap counts, the incidence of PD was not significantly reduced in vineyards planted with highly sensitive varieties (8). Even with a 10-18 foot screen barrier, the number of BGSS flying over the top could still result in significant amounts of PD in an adjacent vineyard.




Figure 4.




Figure 5.

CONCLUSIONS

Nearly 90% of the BGSS trapped in this study were caught on traps at 15 feet or lower. This suggests that barriers could have a significant impact on reducing the numbers of BGSS entering vineyards. However, this may not be enough to have a major impact on reducing the incidence of PD. In addition, results from one tower indicated that BGSS may reside in some trees early in the season. This could allow for higher than normal flight activity, allowing more BGSS to enter vineyards by flying over a barrier. The effectiveness of barriers at reducing the incidence of PD will likely depend upon the nature of the adjacent riparian plant community, its mix of host plant species and the number of tall host trees.

REFERENCES

1. Almeida, R.P.P. and A.H. Purcell. 2003. Transmission of *Xylella fastidiosa* to grapevines by *Homalodisca coagulata* (Hemiptera: Cicadellidae). J. Econ. Ent. 96(2):264-271.
2. Blua, M.J. and D.J.W. Morgan. 2003. Dispersion of *Homalodisca coagulata* (Hemiptera: Cicadellidae), a vector of *Xylella fastidiosa*, into vineyards in southern California. J. Econ. Ent. 96(5):1369-1374.
3. Feil, H., et al. 2000. Effects of temperature on the flight activity of *Graphocephala atropunctata* (Hemiptera: Cicadellidae). Journal of Economic Entomology 93(1): 88-92.
4. Goodwin, P., A. H. Purcell. 1992. Pierce's disease. Grape Pest Management, 2nd Edition. Oakland, University of California, Division of Agriculture and Natural Resources: 76-84.
5. Insley, E., et al. 2000. Riparian vegetation management for Pierce's disease in North Coast California Vineyards. An information manual from the North Coast PD Task Force. 46pp.
6. Purcell, A.H. 1975. Role of the blue-green sharpshooter, *Hordnia circellata*, in the epidemiology of Pierce's disease of grapevines. Env. Ent. 4:745-752.
7. Purcell, A.H. 1976. Seasonal changes in host plant preference of the blue-green sharpshooter *Hordnia circellata*. Pan-Pacific Ent. 52:33-37.
8. Purcell, A.H. 1979. Control of the blue-green sharpshooter and effects of spread of Pierce's disease of grapevines. J. Econ. Ent. 72(6):887-892.
9. Purcell, A.H. 1981. Vector preference and inoculation efficiency as components of resistance to Pierce's disease in European grape cultivars. Phytopath. 71(4):429-435.
10. Weber, E.A. 2004. Evaluation of Blue-green Sharpshooter flight height. Proceedings, Pierce's Disease Research Symposium, San Diego, pp157-159.

FUNDING AGENCIES

Funding for this project was provided by the CDFA Pierce's Disease and Glassy-winged Sharpshooter Board.