

**REALIZED LIFETIME PARASITISM OF GLASSY-WINGED SHARPSHOOTER EGG MASSES
BY *GONATOCERUS ASHMEADI***

Project Leader:

Mark S. Hoddle
Department of Entomology
University of California
Riverside, CA 92521
mhoddle@ucr.edu

Collaborators:

Nic Irvin
Department of Entomology
University of California
Riverside, CA 92521

Robert Luck
Department of Entomology
University of California
Riverside, CA 92521

Reporting Period: The results reported here are from work conducted October 2005 to September 2006.

ABSTRACT

Oosorption and egg maturation results suggested that *Gonatocerus ashmeadi* is a pro-synovigenic species and females mature more eggs during their lifetime. In the absence of hosts, oosorption was initiated on day 7, where the number of reabsorbed eggs increased at a rate of 1-4 eggs per day. In the presence of hosts female *G. ashmeadi* matured 3-27 eggs per day.

INTRODUCTION

The self-introduced *Gonatocerus ashmeadi* (Vickerman et al. 2004) is the key natural enemy of glassy-winged sharpshooter (GWSS; *Homalodisca vitripennis*; formerly *H. coagulata*) egg masses in CA at present (Pilkington et al. 2005). Over summer, parasitism levels of GWSS egg masses and individual eggs in masses by *G. ashmeadi* can approach 100% but parasitism levels of the spring generation of GWSS are substantially lower, and parasitism generally averages ~19-20% (Pilkington et al., 2005; Triapitsyn and Phillips 2000). Naturally occurring populations of *G. ashmeadi* in CA have been augmented with mass reared individuals from populations found in the southeastern U.S.A. and northeastern Mexico which encompasses the home range of GWSS (CDFA 2003).

Substantial laboratory work with *G. ashmeadi* has been conducted in an attempt to understand and parameterize basic aspects of this parasitoid's reproductive biology, and host selection behaviors. Irvin and Hoddle (2005a) have evaluated oviposition preferences of *G. ashmeadi* when presented GWSS eggs of various ages. Interspecific competition between *G. ashmeadi* with *G. triguttatus* and *G. fasciatus* for GWSS egg masses of different ages has been assessed (Irvin and Hoddle 2005b; Irvin et al. 2005) along with factors influencing the sex ratio of offspring (Irvin and Hoddle 2006a). The effect of resource provisioning and nutrient procurement on the longevity of *G. ashmeadi* has also been determined (Irvin and Hoddle 2006b). Furthermore, Pilkington and Hoddle (2006) have assessed laboratory-level fecundity rates of *G. ashmeadi* under different constant temperature regimens.

The GWSS-*Gonatocerus* system has benefited from this intensive laboratory study to generate a basic understanding of factors influencing host selection and parasitism success. The next step that is now required is to test hypotheses generated from lab studies in the field. Field level assessments will help determine the most important aspect of the GWSS biological control program: "How big an impact do individual female *G. ashmeadi* parasitoids have on GWSS population growth via parasitization of eggs?" Addressing this question will allow us to form a much better understanding of the levels of control we can expect from *G. ashmeadi* individually and collectively on GWSS population growth in the field during the spring and summer generations.

OBJECTIVES

To measure real lifetime contributions of individual female *G. ashmeadi* to the parasitism of GWSS egg masses in citrus orchards. Before field assessments can be conducted, laboratory studies will be run to ascertain and verify four critical factors outlined below. Answers to these four critical factors will allow us to develop a composite index that describes the correlative relationship of these four factors that will predict parasitoid age and egg load in the field and to assess the contribution of individual female parasitoids to GWSS suppression under field conditions. The four critical factors are:

1. Determine the relationship between adult female *G. ashmeadi* size as measured by right hind tibia length (HTL) and 24-hr egg load for spring and summer generations (this work was completed and reported in Hoddle et al. 2005).
2. Ascertain the extent to which oosorption occurs, and the length of time without ovipositing that is required to initiate this physiological response if it does occur.
3. Determine whether female parasitoids can mature eggs in excess of those they are born with.
4. Estimate parasitoid age using near infrared spectroscopy (NIRS) (Perez-Mendoza et al. 2002) and develop an alternative measure for comparison by developing a wing deterioration index that estimates parasitoid "age" through visually grading the severity of 'wear and tear' (i.e., numbers of broken setae) of setae on wings (this work was completed and reported in Hoddle et al. 2005).

RESULTS AND DISCUSSION

Hoddle et al. (2005) reported the relationship between *G. ashmeadi* size (tibia length) and <24 hr egg load and described two methods of identifying female age through a wing wear index and using NIRS. Research reported here details oosorption and egg maturation rates for *G. ashmeadi*. These results are preliminary as we are still working on more thorough statistical analyses. With this information we aim to develop a composite index that describes the correlative relationship of the four factors listed above that will predict parasitoid age and egg load in the field and this will allow us to assess the contribution of individual female parasitoids to GWSS suppression under field conditions at time of death.

Oosorption

Gonatocerus parasitoids are generally classified as strictly pro-ovigenic (Jervis and Copland, 1996) where females emerge with a full load of mature eggs and do not mature more eggs as they age (Quicke, 1997). Results obtained so far suggest that *G. ashmeadi* may be partially syn-ovigenic. Completed studies suggest that females emerge with ~30 mature eggs and can mature more over the course of their life time (Figure 1). *G. ashmeadi* females that have access to 50% honey-water but not GWSS eggs developed 4-8 mature eggs per day up to around seven days of age before egg load in females began to decline at a rate of 1-8 eggs per day because of oosorption (Figure 1). In the absence of hosts, females appear to reabsorb mature eggs theoretically enabling them to redirect energy into host seeking and survival, a characteristic of syn-ovigenic species (Jervis et al., 1996). Figure one demonstrates that female *G. ashmeadi* oosorption was initiated on day 7, and the number of reabsorbed eggs increased at a rate of 1-4 eggs per day, to 12 eggs on day 13. In this species, oosorption is obligatory because egg maturing continues in the absence of hosts (Quicke, 1997). However, results also show that the total number of eggs present in *G. ashmeadi* ovaries (potential fecundity) over their lifetime was similar to the predicted <24 hr potential fecundity as estimated from hind tibia length (using data from objective A above). This demonstrates that in the absence of hosts, female *G. ashmeadi* do not mature more eggs than what they emerge with, a characteristic of pro-ovigenic species. In this study, parasitoid age can be converted to physiological age using day-degree estimates (Pilkington and Hoddle, 2006).




Figure 1. The effect of age on the mean number of mature eggs, immature eggs, reabsorbed eggs, potential fecundity and predicted potential fecundity (as predicted from hind tibia length) in ovaries of female *G. ashmeadi* presented with no hosts for 13 days.

Egg Maturation

Figure 2 shows that on day 13, potential fecundity was 77 eggs higher than the predicted <24 hr potential fecundity as estimated from hind tibia length. This suggests that *G. ashmeadi* mature more eggs as they parasitize hosts during their lifetime and indicates that this species is partially syn-ovigenic. Potential fecundity (realized fecundity + eggs present in ovaries) data demonstrates that in the presence of hosts female *G. ashmeadi* matured 3-27 new eggs per day. It is possible that potential fecundity was underestimated in this study because some eggs oviposited by females may be unaccounted for due to superparasitism or early larval death. The potential fecundity of females given hosts for one day after emergence was 26 eggs lower than the predicted <24 hr potential fecundity. This may indicate that 26 eggs were lost due to superparasitism or early larval death.




Figure 2. The number of mature and non-mature eggs present in ovaries, realized fecundity (successful parasitism), potential fecundity (realized fecundity + eggs present in ovaries) and predicted fecundity (estimated from hind tibia length) of female *G. ashmeadi* offered hosts daily for 0-13 days after female emergence.

CONCLUSIONS

Oosorption and egg maturation results suggest that *G. ashmeadi* is a pro-synovigenic species. Females have the ability to mature eggs in excess of those they emerge with over their lifetime. In the absence of hosts, oosorption was initiated on day 7, where the number of reabsorbed eggs increased at a rate of 1-4 eggs per day. In the presence of hosts, female *G. ashmeadi* matured 3-27 new eggs per day. Together with previous data (the relationship between adult female *G. ashmeadi* size and 24-hr egg load, and a wing deterioration index that estimates parasitoid age), these components will be used to develop a composite index that will predict parasitoid age and egg load in the field and help determine how many eggs individual female *G. ashmeadi* parasitize in the field up to the time of death. In 2006 we collected ~20 dead female *G. ashmeadi* from the field using funnel traps loaded with dry ice. Females will be aged and egg load at time of emergence will be estimated from hind tibia length. The egg load at time of death (when oosorption and egg maturation are figured into the model) will allow us to estimate the average number of GWSS eggs females parasitize before dying. These estimates of realized field fecundity will allow us to form a much better understanding of what levels of control individual *G. ashmeadi* in the field are achieving.

REFERENCES

- CDFA. 2003. Pierce's Disease Program Report to the Legislature, May, 2003. California Department of Food and Agriculture.
- Hoddle, M.S. N.A. Irvin, and R. Luck. 2005. Realized lifetime parasitism of glassy-winged sharpshooter egg masses by *Gonatocerus ashmeadi*, pp. 330-333. In: Proceedings, Pierce's Disease Research Symposium, December 5-7, 2004, San Diego, CA.
- Irvin, N. A. and M. S. Hoddle. 2005a. Determination of *Homalodisca coagulata* (Hemiptera: Cicadellidae) egg ages suitable for oviposition by *Gonatocerus ashmeadi*, *Gonatocerus triguttatus*, and *Gonatocerus fasciatus* (Hymenoptera: Mymaridae). *Biological Control* 32: 391-400.
- Irvin, N. A. and M. S. Hoddle. 2005b. The competitive ability of three mymarid egg parasitoids (*Gonatocerus* spp.) for glassy-winged sharpshooter (*Homalodisca coagulata*) eggs. *Biological Control* 34: 204-214.
- Irvin, N. A. and M. S. Hoddle. 2006a. The effect of intraspecific competition on progeny sex ratio in *Gonatocerus* spp. for *Homalodisca coagulata* egg masses: economic implications for mass rearing and biological control. *Biological Control* (in press).
- Irvin, N. A., and M. S. Hoddle. 2006b. Evaluation of floral resources for enhancement of fitness of *Gonatocerus ashmeadi*, an egg parasitoid of the glassy-winged sharpshooter, *Homalodisca vitripennis*. *Biological control* (in press).
- Irvin, N.A., M. S. Hoddle, D. J. W. Morgan, 2006. Competition between *Gonatocerus ashmeadi* and *G. triguttatus* for glassy winged sharpshooter (*Homalodisca coagulata*) egg masses. *Biocontrol Science and Technology* 16 (4): 359-375.
- Jervis, M. A., M. J. W. Copland. 1996. In: Jervis, M. A., Kidd, N. A. (Eds.), *Insect Natural Enemies: Practical approaches to their study and evaluation*. Chapman & Hall, London UK, pp. 63-161.

- Jervis, M. A., N. A. Kidd, , G. E. Heimpel. 1996. Parasitoid adult feeding behaviour and biocontrol – a review. *Biocontrol News and Information* 17(1), 11N-26N.
- Perez-Mendoza, J., F. E. Dowell, A. B. Broce, J. E. Throne, R. A. Wirtz, F. Xie, J. A. Fabrick, and J. E. Baker. 2002. Chronological age-grading of house flies using near infrared spectroscopy. *Journal of Medical Entomology* 39: 499-508.
- Pilkington, L. J., and M. S. Hoddle. 2006. Reproductive and developmental biology of *Gonatocerus ashmeadi* (Hymenoptera: Mymaridae), an egg parasitoid of *Homalodisca coagulata* (Homoptera: Cicadellidae). *Biological Control* (in press).
- Pilkington, L. J., N. A. Irvin, E. A. Boyd, M. S. Hoddle, S. V. Triapitsyn, B. G. Carey, W. A. Jones, and D. J. W. Morgan. 2005. Biological control of glassy-winged sharpshooter in California. *California Agriculture* 59(4), 223-228.
- Quicke, D. L. J. 1997. Parasitic wasps. Chapman and Hall, London.
- Triapitsyn, S. V., P. A. Phillips. 2000. First record of *Gonatocerus triguttatus* (Hymenoptera: Mymaridae) from eggs of *Homalodisca coagulata* (Homoptera: Cicadellidae) with notes on the distribution of the host. *Florida Entomologist* 83, 200-203.
- Vickerman, D.B., M.S. Hoddle, S.V. Triapitsyn, R. Stouthamer. 2004. Species identity of geographically distinct populations of the glassy-winged sharpshooter parasitoid *Gonatocerus ashmeadi*: Morphology, DNA sequences and reproductive compatibility. *Biol. Control* 31(3), 338-345.

FUNDING AGENCIES

Funding for this project was provided by the CDFA Pierce's Disease and Glassy-winged Sharpshooter Board.

DEVELOPING DAY-DEGREE MODELS TO PREDICT SPREAD WITHIN CALIFORNIA OF *GONATOCERUS TRIGUTTATUS* RELEASED FOR GLASSY-WINGED SHARPSHOOTER CONTROL, AND USING DEMOGRAPHIC DATA TO INVESTIGATE PARASITOID SPREAD IN CALIFORNIA

Project Leader:

Mark S. Hoddle
Department of Entomology
University of California
Riverside, CA 92521
mhoddle@ucr.edu

Researcher:

Leigh Pilkington
Department of Entomology
University of California
Riverside, CA 92521

Reporting Period: The results reported here are from work conducted March 2005 to April 2006.

ABSTRACT

The reproductive and developmental biology of *Gonatocerus triguttatus* Girault, a parasitoid of the glassy-winged sharpshooter (GWSS; *Homalodisca vitripennis*; formerly *H. coagulata*), was determined at five constant temperatures in the laboratory; 15; 20; 25; 30; and 33°C. At 25°C, *G. triguttatus* maintained the highest successful parasitism rates with 25.1% of parasitoid larvae surviving to adulthood and lowest parasitism was observed at 15°C with 7.3% of parasitoid larvae surviving to adulthood. Lifetime fecundity was greatest at 25°C and fell sharply as temperature either increased or decreased around 25°C. Temperature had no effect on sex ratio of parasitoid offspring. Mean adult longevity was inversely related to temperature with a maximum of 20.6 days at 15°C to a minimum of four days at 33°C. Developmental rates increased nonlinearly with increasing temperatures. Developmental rate data was fitted with the modified Logan model for oviposition to adult development times across each of the five experimental temperatures. Optimal, lower, and upper lethal, temperature thresholds for *G. triguttatus* were, 30.7°C, 10.4°C and 38.8°C, respectively. The lower developmental threshold estimated with linear regression was 10.57°C, and is very close to the lower temperature threshold estimated by the modified Logan model. Linear regression of developmental rate across all five experimental temperatures indicated that 204 degree-days were required above the minimum threshold of 10.57°C to complete development. Demographic parameters were calculated and pseudo replicates for intrinsic rate of increase (r_m), net reproductive rates (R_o), generation time (T_c), population doubling time (T_d), and finite rate of increase (λ) were generated using the bootstrap method. Mean bootstrap estimates of demographic parameters were compared across temperatures using ANOVA and nonlinear regression.

INTRODUCTION

Gonatocerus triguttatus Girault, (Hymenoptera: Mymaridae) is a solitary endoparasitoid that attacks eggs of sharpshooters in the cicadellid tribe Proconiini. This parasitoid was originally described from specimens reared from an unidentified leafhopper egg mass collected in Trinidad, and has been subsequently reared from glassy-winged sharpshooter (GWSS; *Homalodisca vitripennis*; formerly *H. coagulata*; Hemiptera: Cicadellidae) egg masses collected in Texas and Mexico. This natural enemy has a natural range that includes southeastern U.S.A. and northeastern Mexico where it is associated with GWSS. *Gonatocerus triguttatus* was deliberately imported from Texas U.S.A. and introduced into California U.S.A. in 2002 as part of a classical biological control program against GWSS. Some recoveries from release areas have been made tentatively suggesting *G. triguttatus* may have established perennial populations in California (Pilkington et al., 2005). A thorough understanding and characterization of biological attributes of natural enemies such as degree-day requirements, and intrinsic rates of increase can have multiple practical applications, such as: (1) quantification of the reproductive and developmental biology of candidate natural enemies can assist with predicting potential establishment and population growth of natural enemies introduced into a new area, (2) can aid preliminary evaluation of natural enemies for use potential use in classical biological control, (3) assist with interpretation of natural enemy impact and spread in the field, and (4) provide realistic values for parameters of models investigating incursion risks pertaining to movement of natural enemies into environments beyond those intended for permanent inhabitation. Improved understanding of the basic biology of *G. triguttatus*, a recently released and established natural enemy of GWSS in California, will assist mass-rearing efforts of this parasitoid; optimize timing of inoculative field releases; facilitate better understanding of parasitoid spread and impact on GWSS in various climatic zones in California; and will assist with targeted collecting for biotypes of *G. triguttatus* in the home range of GWSS that may exhibit unique climatic adaptations that current parasitoid populations in California lack.

OBJECTIVES

1. Develop day-degree models for mymarid parasitoids by quantifying the developmental and reproductive biology of *G. triguttatus* at 5 different temperatures (this work has been completed for the principal egg parasitoid of GWSS, *G. ashmeadi*).
2. Use day-degree data from Objective 1 in a Geographic Information Systems approach to predict the geographic range of parasitoids within California and use GIS to map these predictions to known and potential GWSS distributions.