

GRAPE ROOTSTOCK VARIETY INFLUENCE ON PIERCE'S DISEASE SYMPTOMS IN CHARDONNAY

Project Leaders:

Peter Cousins

USDA, ARS

Grape Genetics Research Unit

New York State Agric. Exp. Station

Geneva, New York 14456

peter.cousins@ars.usda.gov

John Goolsby

USDA, ARS

Beneficial Insects Research Unit

Kika de la Garza Subtrop. Ag. Res. Ctr.

Weslaco, Texas 78596

john.goolsby@ars.usda.gov

Reporting Period: The results reported here are from work conducted October 1, 2006 to September 30, 2007.

ABSTRACT

Chardonnay is a *Vitis vinifera* scion variety that is susceptible to Pierce's disease (PD). We are evaluating the effect of rootstock variety on PD symptom expression in Chardonnay grown in an experimental vineyard at Weslaco, Texas with high natural PD pressure and abundant vectors, including glassy-winged sharpshooter. The rootstocks under evaluation are Dog Ridge, Florilush, Tampa, Lenoir, and Freedom. Natural *Xylella fastidiosa* (Xf) infection will be permitted to test the effect of rootstock variety on PD in the Chardonnay scions.

INTRODUCTION

Rootstocks are widely in use in viticulture to manage damage from soil-borne pests and provide adaptation to soils. In citrus (He et al. 2000) and peach (Gould et al. 1991), rootstock variety has been reported to impact expression of Xf diseases in scions. Pierce (1905) reported that rootstock variety affected expression of "California vine disease" (PD) in grape. Grape rootstock trials in Mississippi showed a large effect of rootstock trial on vine longevity in a region recognized for high PD pressure (Loomis 1952, 1965, Magoon and Magness 1937). If grape rootstocks could contribute PD resistance or tolerance to their scions, this would be a major benefit to viticulture in PD prone areas. Elite wine, juice, and table grape varieties could be grown in areas where viticulture is currently restricted to PD resistant and tolerant varieties whose consumer appeal is low.

The Rio Grande Valley is an excellent location for the field evaluation of PD resistant plant germplasm and PD management techniques. Many insect vectors of Xf are native to the region, including the glassy-winged sharpshooter. Susceptible grapevine varieties are infected naturally with Xf in the vineyard and demonstrate characteristic PD symptoms and decline. The Rio Grande Valley is similar to many viticultural regions in California; the region is flat, irrigated, and supports multiple types of crops (citrus, grains, vegetables) in close proximity. The Rio Grande Valley is an ideal test environment due to heavy PD pressure, with abundant vectors and inoculum, in contrast to many other locations, especially California, which demonstrate substantial cycling of PD incidence. The USDA Agricultural Research Service Kika de la Garza Subtropical Agricultural Research Center in Weslaco, Texas is located in the heart of the Rio Grande Valley and provides an ideal experimental location for the evaluation of PD management practices, including rootstock evaluation.

Five rootstocks were chosen for evaluation in this project. Freedom is a complex interspecific hybrid developed as a root-knot nematode resistant rootstock by the USDA ARS, Fresno, California (Clark 1997); its parentage includes *Vitis vinifera*, *V. labrusca*, *V. x champinii*, *V. solonis*, and *V. riparia*. Freedom is widely used in California viticulture. Dog Ridge is a *V. x champinii* selection recognized for its nematode resistance and resistance to PD, but it is rarely used as a rootstock. Lenoir, most probably a *V. aestivalis*/*V. vinifera* hybrid, was used historically as a rootstock and presently is cultivated as a wine grape in PD prone regions (including some parts of Texas) (Galet 1988). Tampa (Mortensen and Stover 1982) includes a *V. aestivalis* selection and the juice grape Niagara (a *V. labrusca* hybrid) in its parentage. Florilush (Mortensen et al. 1994) is a selection from the cross Dog Ridge x Tampa. Both Florilush and Tampa were selected by the University of Florida as PD resistant rootstocks for bunch grapes. PD resistance is necessary for rootstock mothervines to thrive in Florida, so the PD resistance of Florilush and Tampa should not be construed necessarily as contributing to the PD response of the scions.

OBJECTIVE

1. To evaluate the impact of rootstock variety on expression of PD symptoms in naturally infected PD susceptible *Vitis vinifera* scion varieties Chardonnay.

RESULTS AND CONCLUSIONS

Grafted vines of Chardonnay on five rootstocks (Freedom, Tampa, Dog Ridge, Florilush, and Lenoir) were planted at the Kika de la Garza Subtropical Agricultural Research Center in Weslaco, Texas in July, 2006. Evaluation of PD response of the vines began in 2007. Experimental vineyard establishment was good and several vines flowered and fruited in 2007.

Symptoms on leaves were assessed in August 2007 and vines given a numerical score from 0 to 5, with 0 representing no symptoms, 1 = minor symptoms up to 15% of leaves with marginal necrosis (MN), 2 = 15-30% of leaves with MN, 3 = 30-

50% of leaves with MN, 4 = 50-75% of leaves with MN, 5 = over 75% of leaves with MN or vine dead. All vines in the study (total vines = 47) showed MN symptoms. In general, symptom development was minor, with 46 of 47 vines in category MN = 1. Only one vine was assigned to category MN = 2, a vine grafted on the rootstock Tampa.

The rapid development of PD symptoms on the vines in this project following natural inoculation under vineyard conditions reflects the suitability of this location for PD management projects.

REFERENCES

- Clark, J. R. 1997. Grape, pp 248-299. In: The American Society for Horticultural Sciences, ed. The Brooks and Olmo Register of Fruit and Nut Varieties. ASHS Press, Alexandria, Virginia.
- Galet, P. 1988. Cépages et Vignobles de France--Tome I: Les Vignes Américaines. Charles Déhan, Montpellier, France.
- Gould, A. B., W. J. French, J. H. Aldrich, B. V. Brodbeck, R. F. Mizell III, and P. C. Andersen. 1991. Rootstock influence on occurrence of *Homalodisca coagulata*, peach xylem fluid amino acids, and concentrations of *Xylella fastidiosa*. Plant Dis. 75:767-770.
- He, C. X., Li, W. B., Ayres, A. J., Hartung, J. S., Miranda, V. S., and Teixeira, D. C. 2000. Distribution of *Xylella fastidiosa* in citrus rootstocks and transmission of citrus variegated chlorosis between sweet orange plants through natural root grafts. Plant Dis. 84:622-626.
- Loomis, N. H. 1952. Effect of fourteen rootstocks on yield, vigor, and longevity of twelve varieties of grapes at Meridian, Mississippi. Proceedings of the American Society for Horticultural Science. 52:125-132.
- Loomis, N. H. 1965. Further trials of grape rootstocks in Mississippi. Proceedings of the American Society for Horticultural Science 86:326-328.
- Magoon, C. A. and J. R. Magness. 1937. Investigations on the adaptability of grape root stocks to Gulf Coast conditions. Proceedings of the American Society for Horticultural Science 35:466-470.
- Mortensen, J. A., J. W. Harris, and D. L. Hopkins. 1994. 'Florilush': a bunch grape rootstock for Florida. HortScience 29:1375-1376.
- Mortensen, J. A. and L. H. Stover. 1982. Tampa—a new bunch grape rootstock. Agricultural Experiment Stations, Institute of Food and Agricultural Sciences, University of Florida, Gainesville. Circular S-295.
- Pierce, N. B. 1905. The Vineyard: Mr. Pierce and the Lenoir. Pacific Rural Press, 69:79.

FUNDING AGENCIES

Funding for this project was provided by the USDA Agricultural Research Service.

Additional Note: Special thanks to Duarte Nursery, Hughson, California for custom propagation of the experimental grapevines and to Professor M. Andrew Walker, Department of Viticulture and Enology, University of California, Davis, for Lenoir cuttings.