

SEASONAL TRANSMISSION OF *XYLELLA FASTIDIOSA* BY THE GLASSY-WINGED SHARPSHOOTER FROM GRAPEVINES INFECTED FOR VARIOUS LENGTHS OF TIME

Project Leader:

Thomas M. Perring
Department of Entomology
University of California
Riverside, CA 92521
thomas.perring@ucr.edu

Cooperators:

Charles A. Farrar
Department of Entomology
University of California
Riverside, CA 92521
charles.farrar@ucr.edu

Matthew J. Blua
Department of Entomology
University of California
Riverside, CA 92521
matthew.blua@ucr.edu

Reporting Period: The results reported here are from work conducted July 2007 through September 2007.

ABSTRACT

This study was designed to define specific orientation and feeding behaviors by the glassy-winged sharpshooter (GWSS) in association with grapevines that influence vine-to-vine spread of *Xylella fastidiosa* (*Xf*), the Pierce's disease bacterium. Thus far, we conducted experiments in which we made hourly observations on the location of individual GWSS adults given access to mature tissue and a terminal growing tip on the same cane. Although GWSS were observed on all tissues, both males and females preferred immature tissues to mature tissues on Cabernet Sauvignon and Chardonnay grapevines, yet sharpshooters moved frequently throughout the day. Immature stems and leaves were consistently the most used tissue, while immature petioles and mature stems and petioles were the least used.

INTRODUCTION

Pierce's disease (PD), a disease of grapes caused by the bacteria, *Xylella fastidiosa* (*Xf*) Wells et al., was described in California in the 1880s during an epidemic in Orange County (Pierce 1882). A second epidemic occurred in Tulare County in the 1930s (Hewitt et al. 1949), and until the mid-1990s, it was considered only a minor problem in vineyards close to riparian areas. In the early 1990s a new vector, the GWSS, *Homalodisca vitripennis* (Germar) (formerly *Homalodisca coagulata* Say), was introduced into the state (Sorenson and Gill 1996), and became associated with a devastating epidemic of PD in the Temecula Valley. Since 1994, at least 1,500 acres of vineyards have been lost to the disease in California; in the Temecula Valley alone, losses have been estimated at \$13 million (Wine Institute 2002). The GWSS has different feeding and dispersal capabilities than native insect sharpshooter vectors and these attributes are thought to have contributed to the increased number of PD-infected grapevines in California (Almeida et al. 2005a, Blua et al. 1999, Redak et al. 2004). Like other insect-borne plant pathogen systems, there are two potential types of pathogen spread: primary or secondary spread. Primary spread occurs when the pathogen is obtained by the vector from sources outside the crop and transported and inoculated into the crop. Secondary spread occurs when the vector acquires the pathogen from infected vines in the vineyard, and subsequently inoculates healthy vines within the same vineyard (i.e. vine to vine spread). It is thought that *Xf* spread with native California vectors was the result of primary spread, but that rapid spread by GWSS may be the consequence of primary and secondary spread (Almeida et al. 2005a, Hill 2006). Consequently, the best strategy for managing PD spread by GWSS is to control the vector. This strategy, which is simple in concept, yet difficult and costly to implement over large geographic areas, is exactly the approach of the area-wide management program being implemented by USDA, CDFA, County Agricultural Commissioner's Offices, and the University of California (for example Toscano and Gispert 2005, Stone-Smith et al. 2005). This strategy has been so effective at slowing the spread of PD throughout the state that the "crisis" phase of the PD-GWSS problem appears to have passed. Unknown, of course, is what will happen should the area-wide management program be reduced. Due to this uncertainty, researchers continue to search for alternative solutions, both short and long term. Understanding details of primary and secondary spread of *Xf* by GWSS can assist in the development of alternatives to the area-wide management program. For example, to reduce primary spread, efforts must focus on reducing bacteria-carrying GWSS from entering healthy vineyards, through continued area-wide or local treatment programs outside the vineyard, barriers, trap crops, and/or removal of pathogen sources outside the vineyard. Reduction of secondary spread can be accomplished by in-field control of GWSS, finding and roguing infected vines in the vineyard (Varela et al. 2001), and/or minimizing acquisition from infected vines and transmission to healthy vines.

The relationship among time of inoculation, location of inoculation, and disease progression in the vine likely plays a role in determining whether disease becomes chronic and when a vine becomes a source plant for additional spread. When another PD vector, the blue-green sharpshooter, *Graphocephala atropunctata*, infected grapevines early in the season, more persistent infections resulted than from later season infection (Purcell 1981). A potential difference between blue-green sharpshooter transmission and GWSS transmission is that the former is known to prefer feeding at the tips of canes (Purcell 1976), whereas the latter has been reported to feed on older plant parts. Almeida et al. (2005b) demonstrated that GWSS could even transmit *Xf* to dormant vines in the field. However acquisitions from dormant vines in the field were negative. Whether these transmissions and acquisitions are important to disease spread depends on GWSS feeding preferences during the winter months when the vines are dormant. Similarly, it is possible that infection at certain times of the season may not become systemic because infection is pruned out at end of year, or environmental conditions limit bacterial spread (Feil and Purcell 2001, Feil et al. 2003, Hill 2006).

OBJECTIVES

While this progress report addresses only the first objective, the objectives of the project are:

1. Document GWSS feeding preference, through the growing season, on established Cabernet Sauvignon and Chardonnay grapevines that either are healthy or have been infected with *Xf* for 2, 3, or 4 years.
2. Evaluate the acquisition by GWSS, through the growing season, from established Cabernet Sauvignon and Chardonnay grapevines that either are healthy or have been infected with *Xf* for 2, 3, or 4 years and determine the subsequent transmission from these acquisitions.
3. Determine the relationship between *Xf* inoculation by GWSS at different times of the year and the development of the vine as a source for further acquisition by GWSS.

RESULTS


We placed GWSS adults individually in observation cages fabricated from acetate cylinders (25cm x 17cm diameter) with organdy sleeves attached to the ends. The cage was placed over the base of a Cabernet Sauvignon or Chardonnay grape vine cane with the cane terminal looped back into it. The ends of the observation cage were sealed giving each GWSS access to immature and mature stems, petioles, and leaves inside. The grapevines were from a mixed field-grown vineyard at the University of California in Riverside that was covered with 60% shade-cloth to protect them from PD. We made hourly observations from 700 to 1800 hours over three days to determine which tissue each GWSS was on. This experiment was conducted in late August (Experiment 1) and again in mid September 2007 (Experiment 2).

In both experiments, GWSS were found on the cage in 12 to 16% of our observations (Figure 1). When on the cane, GWSS were observed on all tissue types, and although significant differences in tissues used occurred among varieties and genders in both experiments, GWSS did not show a substantial preference for a specific tissue consistently. However, they were found more frequently on immature stems, petioles, and leaves (46-67% of our observations) than mature tissue (21-37%) whether they were enclosed on Cabernet Sauvignon or Chardonnay vines, or were males or females (Figure 1). In general, GWSS were most likely to be found on immature stems and leaves, and least likely to be found on immature petioles and mature stems and petioles (Figure 1). Importantly, averages of 34% and 36% of experimental GWSS across gender in experiments 1 and 2, respectively, switched from one tissue to another each hour on Cabernet Sauvignon. Hourly averages of 30% and 37% of GWSS on Chardonnay switched tissues in experiments 1 and 2, respectively. Averages of 34% and 39% of females switched tissues each hour, and averages of 30% and 34% of males switched tissues each hour in experiments 1 and 2, respectively.

CONCLUSIONS

Vine to vine spread of *Xf* by GWSS has been hypothesized as a critical component of devastating PD epidemics that occurred in Temecula and in the General Beale area of Kern County. A fundamental understanding of this type of spread can lead to strategies insuring that epidemics of these proportions do not occur elsewhere. GWSS landing and feeding behavior, and tissue feeding capacity combine with grapevine phenology, and *Xf* phenology to make vine to vine spread possible. Particularly important is the tendency for GWSS to move frequently in grapevines, as shown in this study, and their characteristic short hopping flights (Turner and Pollard 1959) that would maximize within-vineyard spread of *Xf*. Increased movement by GWSS in search for optimal host tissue would increase the chance of contact with infected and healthy grapevines alike. Our future studies will compare GWSS feeding in no-choice circumstances, and GWSS flight orientation and behavior when confronted with PD-infected and healthy grapevines. We will also examine the relationship between *Xf* inoculation by GWSS at different times of the year and the development of the vine as a source for further acquisition by GWSS.

Experiment 1



Experiment 2




Figure 1. GWSS preference on field-grown Cabernet Sauvignon and Chardonnay grapevines in choice experiments conducted on 29 August 2007 (Experiment 1) and 11 September 2007 (Experiment 2). Bars represent average proportions of GWSS (\pm SE) observed on various tissue types. Different letters above bars represent statistically significant differences among means at $p = 0.05$ (ANOVA followed by t-tests for mean separation).

REFERENCES

- Almeida, R. P. P., M. J. Blua, J. R. S. Lopes, and A. H. Purcell. 2005a. Vector transmission of *Xylella fastidiosa*: applying fundamental knowledge to generate disease management strategies. *Ann. Ent. Soc. Am.* 98: 775-786.
- Almeida, R. P. P., C. Wistrom, B. L. Hill, J. Hashim, and A. H. Purcell. 2005b. Vector transmission of *Xylella fastidiosa* to dormant grape. *Plant Dis.* 89: 419-424.
- Blua, M. J., P. A. Phillips, and R. A. Redak. 1999. A new sharpshooter threatens both crops and ornamentals. *Calif. Agric.* 53: 22-25.

- Feil, H., W.S. Feil, and A.H. Purcell. 2003. Effects of date of inoculation on the within-plant movement of *Xylella fastidiosa* and persistence of Pierce's disease within field grapevines. *Phytopathology*. 93: 244-251.
- Feil, H., and A.H. Purcell. 2001. Temperature-dependent growth and survival of *Xylella fastidiosa* in vitro and in potted grapevines. *Plant Dis.* 85: 1230-1234.
- Hewitt, W. B., N.W. Frazier, J.H. Freitag, and A.J. Winkler. 1949. Pierce's disease investigations. *Hilgardia* 19:207-264.
- Hill, B.L. 2006. The effect of dormant season survival of *Xylella fastidiosa* in grapevines on Pierce's disease epidemics in California. Pp. 276-279 in T. Esser (ed.) Symposium Proc. of the 2006 Pierces Disease Research Symposium. Nov. 2006, San Diego, CA. California Department of Food and Agriculture, Sacramento, CA.
- Pierce, N. B. 1882. The California vine disease. U. S. Div. Vegetable Physiology and Pathology. Bulletin No. 2.
- Purcell, A. H. 1976. Seasonal changes in host plant preference of the blue-green sharpshooter *Hordnia circellata*. *Pan-Pacific Entomol.* 52:33-37
- Purcell, A. H. 1981. Vector preference and inoculation efficiency as components of resistance to Pierce's disease in European grape cultivars. *Phytopathology* 71:429-435.
- Redak, R. A., A. H. Purcell, J. R. S. Lopes, M. J. Blua, R. F. Mizell, and P. C. Andersen. 2004. The biology of xylem fluid-feeding insect vectors of *Xylella fastidiosa* and their relation to disease epidemiology. *Annu. Rev. Entomol.* 49: 243-270.
- Sorenson, J. T. and R. J. Gill. 1996. A range extension of *Homalodisca coagulata* (Say) (Hemiptera: Clypeorrhyncha: Cicadellidae) to Southern California. *Pan-Pac. Entomol.* 72: 160-161.
- Stone-Smith, B., J. Stewart-Leslie, G. Kunkel, S. Hardy, B. Appleby, D. Borges, D. Haines, J. Churchill. 2005. The area-wide pest management of glassy-winged sharpshooter in Tulare County, pp. 376-379, In M. A. Tariq, P. Blincoe, M. Mochel, S. Oswalt, and T. Esser (eds.), Proceedings of Pierce's disease research symposium 2005, San Diego, CA.
- Toscano, N.C. and C. Gispert. 2005. Riverside County glassy-winged sharpshooter area-wide management program in the Coachella and Temecula valleys. Pp. 384-387 In M. A. Tariq, P. Blincoe, M. Mochel, S. Oswalt, and T. Esser (eds.), Proceedings of Pierce's disease research symposium 2005, San Diego, CA.
- Turner, W.F., and H.N. Pollard. 1959. Life histories and behavior of five insect vectors of phony peach disease. U.S. Dept. Agric. Tech. Bull. 1188.
- Varela, L. G., R. J. Smith, and P. A. Phillips. 2001. Pierce's Disease. University of California, Agriculture and Natural Resources Publication No. 21600. Wine Institute. 2002. Pierce's disease update.
- Wine Institute. 2002. Pierce's disease update.
http://www.wineinstitute.org/communications/pierces_disease/pierces_disease_update.htm

FUNDING AGENCIES

Funding for this project was provided by the CDFA Pierce's Disease and Glassy-winged Sharpshooter Board.