

**Project Title: Identification of factors mediating cold therapy of *Xylella fastidiosa*-infected grapevines.**

**Principle Investigator and Cooperator:** Bruce C. Kirkpatrick and Melody M. Meyer.  
University of California, Davis.

**Objective 1: Develop an experimental, growth chamber temperature regime that can consistently cure Pierce's Disease affected grapevines without causing unacceptable plant mortality.**

Methods were identical to those used in the 2005-2006 season. Pinot Noir (PN, susceptible to PD) and Cabernet Sauvignon (CS, moderately resistant to PD) grapevines grafted onto 101-14 rootstock were inoculated with *X. fastidiosa* and then exposed to field conditions and cold chamber conditions to define the temperature threshold that induces the cold curing phenomenon and to identify the physiological response within the plant that mediates PD recovery.

**Field Experiment Results:**

The results for our field plots showed lower disease ratings and higher curing rates in the colder temperature treatments (Table 1). This is consistent with the results we obtained in previous field seasons. This information will be used to make a model that will help predict what areas are most likely to experience cold curing and not expected to have PD problems.

**Table 1: 2006-2007 Field Results**

<b>PD-free vines<sup>x</sup>/ grapevines that survived until 2007</b>	<b>Davis</b>	<b>Hopland</b>	<b>McLaughlin</b>	<b>Foresthill</b>
Pinot Noir	3/11	9/11	8/11	8/11
Cabernet Sauvignon	1/11	8/11	7/11	10/11

<b>Mean Disease Ratings after cold treatments<sup>y</sup></b>	<b>Davis</b>	<b>Hopland</b>	<b>McLaughlin</b>	<b>Foresthill</b>
Pinot Noir	1.6	0.6	1.0	0.3
Cabernet Sauvignon	2.3	1.0	1.1	0.1

<b>Vine Mortality<sup>z</sup></b>	<b>Davis</b>	<b>Hopland</b>	<b>McLaughlin</b>	<b>Foresthill</b>
Pinot Noir	0%	0%	0%	0%
Cabernet Sauvignon	0%	0%	0%	0%

<sup>x</sup>. Vines that were inoculated with *X. fastidiosa* and tested (+) by IC-PCR in Fall 2006 that no longer tested (+) in Fall 2007.

<sup>y</sup>. Disease ratings for Pierce's Disease 0-5 (no disease to most severe disease).

<sup>z</sup>. Total mortality of inoculated and control vines.

**Cold Room Experimental Results:**

The results of the cold room experiments showed disease recovery and mortality trends that were similar to the field plots. The coldest treatments had the highest rate of recovery from PD, but also the highest mortality (Table 2).

Table 2: 2006-2007 Cold Chamber Results

PD-free vines <sup>x</sup> / inoculated grapevines that survived until 2007	+5°C	2.2°C	0°C	-5°C
Pinot Noir	3/9	3/10	4/10	6/6
Cabernet Sauvignon	0/9	4/10	5/10	7/7

Mean Disease Ratings after cold treatments <sup>y</sup>	+5°C	2.2°C	0°C	-5°C
Pinot Noir	1.4	2.1	1.1	0.0
Cabernet Sauvignon	2.1	2.2	0.6	0.0

Vine Mortality <sup>z</sup>	+5°C	2.2°C	0°C	-5°C
Pinot Noir	10%	0%	0%	40%
Cabernet Sauvignon	10%	0%	0%	30%

<sup>x</sup>. Vines that were inoculated with *X. fastidiosa* and tested (+) by IC-PCR in Fall 2006 that no longer tested (+) in Fall 2007.

<sup>y</sup>. Disease ratings for Pierce's Disease 0-5 (no disease to most severe disease).

<sup>z</sup>. Total mortality of inoculated and control vines.

**Objective 2: Analyze chemical changes such as pH, osmolarity, total organic acids, proteins and other constituents that occur in the xylem sap of cold-treated versus non-treated susceptible and less susceptible *Vitis vinifera* varieties.**

Field grown and growth chamber plants were prepared as described in Objective 1. Xylem sap was extracted from vines from each field location and cold chamber treatment using the pressure bomb. The samples were then tested for potential changes in pH, osmolarity, protein profiles, total sugars, and calcium and magnesium concentrations in xylem sap.

Xylem sap protein profiles were analyzed for the 2005-2007 samples. The sap proteins were concentrated with acetone precipitation and then electrophoresing the proteins in a 12% Tris-HCl 1-dimensional polyacrylamide gel (PAGE). Protein profiles of the PAGE gels were compared for each treatment. Unique protein bands that were found in the cold treated plants were cut from the gel, and end terminally sequenced by the UCD Molecular Structure Facility.

**Field Experiment Results:**

ABA concentrations in the spring xylem sap collections were the lowest in the coldest field locations (Table 3). ABA levels were higher in the late winter sap collections than in the spring collections for the field locations.

**Cold Chamber Experiment Results:**

The results of the cold chamber experiments show that pH of CS xylem sap was significantly higher than PN sap overall (Table 4). Sugar and select ion concentration analysis of CS grapevines showed greater amounts of glucose and fructose in -5°C cold chamber vines, whereas Ca<sup>+</sup> levels were greater in the warmest treatments (Figure 1). Osmolarity was greatest in the coldest treatments and decreased with increasing temperature. Conversely, in PN grapevines, glucose and fructose levels were the lowest in the coldest treatments (Figure 2). Ca<sup>+</sup> levels showed a similar trend with CS vines, with increased Ca<sup>+</sup> levels in the warmer temperature treatments. Temperature appeared to have a less direct effect on osmolarity in Pinot Noir

grapevines. ABA concentrations in the spring xylem sap collections were the lowest in the coldest cold chambers (Table 4).

While most of the proteins were similar for the various temperatures, a few unique proteins were found in the cold stressed and/or *X. fastidiosa*-inoculated plants and these proteins were end terminally sequenced by the UCD Molecular Structure Facility. Sequencing results of xylem proteins from cold-treated vines showed proteins that had high sequence homology with stress proteins that are produced by Cabernet Sauvignon berries under water deficit stress conditions, proteins that are similar to proteins produced in Pinot Noir roots, trypsin inhibitors and a thaumatin-like protein (Figure 3).

Table 3: 2006-2007 Mean osmolarity, pH, and abscisic acid concentration of xylem sap from grapevines growing at 4 field locations around California.

			Davis		Hopland		McLaughlin		Foresthill	
			1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>	1 <sup>st</sup>	2 <sup>nd</sup>
<b>pH</b>	Pinot Noir	Control	5.36	6.09	4.92	6.07	6.06	5.80	6.17	6.10
		Inoculated	5.88	6.34	5.46	5.76	6.00	6.00	5.92	6.08
	Cabernet Sauvignon	Control	5.77	6.35	5.14	5.74	6.41	5.04	6.21	6.27
		Inoculated	6.00	6.42	4.99	5.81	6.19	5.76	6.24	6.23

<b>Osmolarity</b>	Pinot Noir	Control	78.1	61.6	80.7	54.4	118.7	48.0	179.5	67.5
		Inoculated	104.5	43.1	78.7	79.9	76.0	31.4	67.2	57.8
<b>mmol/kg</b>	Cabernet Sauvignon	Control	95.5	34.6	74.4	77.1	78.9	24.5	134.7	61.1
		Inoculated	102.1	35.8	72.1	76.5	84.3	25.0	86.6	53.1

<b>Absciscic Acid</b>	Pinot Noir	Control	387	722	364	264	350	354	384	504
		Inoculated	536	833	238	388	234	150	418	319
<b>pmol/mL</b>	Cabernet Sauvignon	Control	320	558	252	282	464	390	584	554
		Inoculated	254	278	452	500	530	232	304	273

Table 4: Osmolarity, pH, and ABA concentrations of xylem sap from grapevines treated with 4 different cold temperature regimes in growth chambers 2006-2007.

			-5°C	+0°C	+2.2°C	+5°C
<b>pH</b>	Pinot Noir	Control	6.36	5.88	5.38	5.86
		Inoculated	5.66	5.68	5.52	6.00
	Cabernet Sauvignon	Control	5.66	5.89	5.82	6.31
		Inoculated	5.78	5.83	5.85	6.30

<b>Osmolarity</b>	Pinot Noir	Control	32.3	48.4	39.2	100.0
		Inoculated	37.2	56.9	39.9	69.9
<b>mmol/kg</b>	Cabernet Sauvignon	Control	37.9	57.4	46.6	67.5
		Inoculated	25.3	26.5	32.4	56.3

<b>Absciscic Acid</b>	Pinot Noir	Control	804	254	450	570
		Inoculated	375	201	210	450
<b>pmol/mL</b>	Cabernet Sauvignon	Control	652	362	166	358
		Inoculated	274	396	488	246

Figure 1: Fructose, glucose, Ca<sup>+</sup> and Mg<sup>+</sup> concentrations from cold room treated Cabernet Sauvignon grapevines plotted against osmolarity of xylem sap.




Figure 2: Sugar (fructose and glucose) and ion (Ca<sup>+</sup> and Mg<sup>+</sup>) concentrations in xylem sap from cold treated Pinot Noir grapevines compared to osmolarity.





Figure 3: Protein profile of grapevine xylem sap. 150 uL of xylem sap was precipitated with cold acetone. Proteins were resuspended in 30 uL of SDS-loading buffer and loaded in to a BioRad 12% Tris-HCl gel



**Objective 3: Assess the viability of cultured *X. fastidiosa* cells growing in media with varying pH and osmolarity and cells exposed to xylem sap extracted from cold- and non-treated grapevines.**

The solutions used for these viability experiments included: water, extracted *V. vinifera* ('Pinot Noir' and 'Cabernet Sauvignon' varieties) xylem sap, PD3, HEPES, sodium and potassium phosphate buffers. All buffers and media were adjusted to pH 6.8. *X. fastidiosa* cells suspended in the various buffers and media were exposed to various temperatures (28°C, 5°C, 2.2°C, 0°C, -5°C, -10°C and -20°C). Potassium phosphate buffer at various pH values (5.0-6.8) was also used to determine the effects of pH on the survival of *X. fastidiosa*. The 10<sup>7</sup> concentration of *X. fastidiosa* cells was determined using a spectrophotometer (Thermo Spectronic; Rochester, NY USA) and once the desired solution was made, portions of the solutions were plated and counted seven days post plating to confirm viable counts.

**Results:**

The results of these experiments were reported in detail in the 2007 progress report. To summarize the results, these experiments indicate that *X. fastidiosa* can survive at 28°C in most media except water. The mortality rate was the lowest in PD3 medium in the 5°C and 2.2°C temperature treatment. The deionized water treatment had the highest mortality rate followed by potassium phosphate at pH 6.2. The highest survival at 0°C occurred with PD3 media and in xylem sap collected from grapevines growing in a cold climate (Placer County, CA). Survival was the lowest in deionized water and potassium phosphate at the pH of 6.2. These experiments indicate that *X. fastidiosa* can survive at -5°C in all buffers at pH 6.8, media and xylem sap for at least 4 days. No cultivable *X. fastidiosa* was recovered from any of the media, buffers or xylem sap after 24 hours at -10°C or at -20°C.

**Objective 4: Determine the effect of treating PD-affected grapevines with cold-induced plant growth regulators, such as abscisic acid (ABA), as a possible therapy for PD.**

We contacted Valent Bioscience Corporation who has an active research and development program on the use of ABA on agricultural crops. In November of 2005, 2006, and 2007, healthy and *X. fastidiosa*-inoculated Cabernet Sauvignon and Pinot Noir vines prepared described in Objective 1 were sprayed with solutions of ABA in the fall.

**2006-2007 Results:**

To evaluate the reproducibility of the 2005-2006 results a second ABA spray trial was conducted in the 2006-2007 season. In 2006 there were 4 treatments with *X. fastidiosa*-infected vines and healthy controls:

- Control: 16 Pinot/ 16 Cabernet plants sprayed with water
- 2000ppm spray: 16 Pinot/ 16 Cabernet plants sprayed with VBC-30054
- 100 ppm drench: 16 Pinot/ 16 Cabernet plants drenched with VBC-30054
- 10 ppm drench: 16 Pinot/ 16 Cabernet plants drenched with VBC-30030


To determine effectiveness of ABA and synthetic ABA treatments on *X. fastidiosa*-infected grapevines, the vines were evaluated for PD symptoms and tested by IC-PCR in the late summer.

Our applications of ABA in the 2005-2006 season appeared to have a curing effect in PD-infected grapevines. ABA application appeared to be the most effective when VBC-30030 was applied as a drench, but some of the other forms and concentrations of ABA seem to also

have some curing effect. For this first application in 2005-2006 there was no rain until a week following the application.

In 2006-2007 this experiment was replicated with some modifications to the treatments as seen above. Curing rates were not the same as what we saw in the 2005-2006 treatments. The only treatment that seemed to have more curing than the control treatment was the VBC-30030 drench in Pinot Noir grapevines, but the curing rate was not as high as in the 2005-2006 season (Figure 4). This could possibly due to a rain event that occurred a few hours after the ABA application, possibly diluting, washing off, or leaching out the applied ABA.

**Figure 4: Percent curing of *Xylella fastidiosa* infected grapevines in the 2006-2007 experiment.**



#### **2007-2008:**

To evaluate the reproducibility of the 2005-2006 results a third ABA spray trial was conducted in the 2007-2008 season. In 2007 there were 4 treatments with *X. fastidiosa*- infected vines and healthy controls:

- Control: 16 Pinot/ 16 Cabernet/ 16 Chardonnay plants sprayed with water
- 2000ppm spray: 16 Pinot/ 16 Cabernet /16 Chardonnay plants sprayed with VBC-30054
- 100ppm spray: 16 Pinot/ 16 Cabernet/16 Chardonnay plants sprayed with VBC-30030
- 100 ppm drench: 16 Pinot/ 16 Cabernet/16 Chardonnay plants drenched with VBC-30054
- 10 ppm drench: 16 Pinot/ 16 Cabernet/16 Chardonnay plants drenched with VBC-30030

The xylem sap of the grapevines was extracted using a pressure bomb four days after the application of the ABA treatments. To examine the proteins produced when grapevines are exposed to ABA, protein profiles were made of each treatment. The 150 ul of xylem sap was precipitated with cold acetone to concentrate the proteins. The proteins were resuspended in 30 uL of SDS-loading buffer and electrophoresed in a BioRad 12% Tris-HCl gel. Some of the proteins sequences in the ABA treated vines are similar to those found in our cold treated vines.

#### **Publications or reports resulting from the project:**

2007 Pierces Disease Research Symposium Report; 2006 Pierces Disease Research Symposium Report; 2005 Pierces Disease Research Symposium Report; 2004 Pierces Disease Research Symposium Report.

#### **Presentations on Research:**

2007 Pierces Disease Research Symposium; 2007 American Phytopathological Society Meeting; 2007 National Viticulture Research Conference; 2006 Pierces Disease Research Symposium; 2006 American Phytopathological Society Meeting; 2005 Pierces Disease Research Symposium; 2004 Pierces Disease Research Symposium.

**Research Relevance Statement:**

PD is currently found in many regions of California and the southern United States. One factor that has been shown to be associated with the observed limited geographical distribution of PD in North America is the severity of winter temperatures in those regions. For example, PD does not occur in New York, the Pacific Northwest or at high altitudes in South Carolina, Texas and California where the winter temperatures on average drop below zero degrees Celsius (Hopkins & Purcell, 2002). Purcell (1977, 1980) and Feil's (2002) research suggested that some factor(s) expressed in the intact plants helps eliminate *X. fastidiosa* from grapevines.

To elucidate the mechanism(s) of the "cold curing" phenomenon, it is necessary to determine the cold curing temperature threshold that maximizes PD recovery and minimizes vine mortality. This research should also allow us to generate projection maps to determine if vineyards in cold boundary areas (i.e., foothills of the Sierra and northern-most California) are at risk for developing PD. The information obtained from these experiments will facilitate basic research on mechanisms causing cold therapy and provide data that could be used by grape growers for risk assessment and management purposes.

It has been well documented that xylem sap contains many metabolites such as mono- and disaccharides, organic acids, plant growth regulators and other organic compounds (Andersen et al., 1989; Bollard, 1960; Pate, 1976; Wormall, 1924; reviewed by Seyedbagheri & Fallahi, 1994). Though it is well known that these compounds are in the sap, little is known about the effects of cold temperatures on the synthesis or quantity of these compounds in sap. Also, little is known about the effect of cold temperatures on factors such as pH and osmolarity of xylem sap and how these factors may be contributing to the cold curing phenomenon. Assessing the effect of pH and osmolarity on the viability of *X. fastidiosa* cells *in vitro*, could provide insight into the factors that contribute to the cold curing phenomenon.

Previous research has shown that herbaceous and woody plants exposed to sub-lethal cold conditions have significantly elevated levels of plant hormones, such as abscisic acid (ABA), which induce the synthesis of a number of cold shock proteins (Guy, 1990; Bravo, et al., 1998; Thomashow, 1998). Kuwabara et al. (2002) elicited cold-shock proteins at 23°C in winter wheat using an exogenously applied 100ppm ABA solution. ABA treated plants elicited proteins that were able to inhibit fungal growth when exposed to exogenous applications of ABA. The application of ABA could lead to a potentially novel approach for managing Pierce's Disease.

**Lay Summary of Current Years Results:**

The results of our field and cold chamber experiments show lower disease ratings and higher curing rates in the colder temperature treatments. The coldest treatments had the highest rate of recovery from PD, but also the highest grapevine mortality.

The analysis of the biochemical factors in sap revealed some interesting results. For the cold chamber experiments the pH of CS xylem sap was significantly higher than PN sap overall. Sugar and select ion concentration analysis of CS grapevines showed greater amounts of glucose and fructose in -5°C cold chamber vines, whereas Ca<sup>+</sup> levels were greater in the warmest treatments. Osmolarity was greatest in the coldest treatments and decreased with increasing temperature. Conversely, in PN grapevines, glucose and fructose levels were the lowest in the coldest treatments. Interestingly, the osmolarity of PD3 media is 113 mmol/kg, whereas the osmolarity of xylem sap, 25-45 mmol/kg. Ca<sup>+</sup> levels showed a similar trend with CS vines, with increased Ca<sup>+</sup> levels in the warmer temperature treatments. Temperature appeared to have a less direct effect on osmolarity in Pinot Noir grapevines.

ABA concentrations in the spring xylem sap collections were the lowest in the coldest field locations and coldest cold chambers. ABA levels were higher in the late winter sap collections than in the spring collections for the field locations.

PAGE protein profile analysis showed that most of the proteins we found were similar for the various temperatures, but a few unique proteins were found in the cold stressed and/or *X. fastidiosa*-inoculated plants. Sequencing results of xylem proteins from cold-treated vines showed proteins that are similar to stress proteins that are produced by Cabernet Sauvignon berries under water deficit stress conditions, proteins that are similar to proteins produced in Pinot Noir roots, and trypsin inhibitors. The protein similarity that is most interesting is a thaumatin-like protein, which has been reported to have anti-microbial activity, that appears to be produced in greater quantities under the coldest conditions. We plan to assess any anti-*X. fastidiosa* properties of this protein in the future.

The *in vitro* culture experiments indicate that *X. fastidiosa* can survive at 28°C in most media except water. At 28°C the survival rate was the highest in PD3 media followed by potassium phosphate at pH 6.8, sodium phosphate, and xylem sap. At the coldest temperatures, the highest survival at 0°C occurred with PD3 media and in xylem sap collected from grapevines growing in a cold climate (Placer County, CA), whereas survival was the lowest in deionized water and potassium phosphate at pH 6.2. Interestingly, *X. fastidiosa* can survive at -5°C in all buffers at pH 6.8, media and xylem sap for at least 4 days. No cultivable *X. fastidiosa* was recovered from any of the media, buffers or xylem sap after 24 hours at -10°C or at -20°C. *X. fastidiosa* in potassium phosphate buffers with pH values at 5.0, 5.4 and 5.8 died rapidly at all temperatures.

The results the ABA application experiments in the 2005-2006 season indicate that ABA appears to have a curing effect when applied as a drench. The synthetic ABA had the most interesting result with 100% curing in Pinot Noir vines. Disease ratings for both drench treatments decreased or were eliminated after application of drench treatments.

In 2006-2007 this experiment was replicated with some modifications to the 2005-2006 treatments. Curing rates were not as high as what we saw in the 2005-2006 treatments. The only treatment that seemed to have more curing than the control treatment was the VBC-30030 drench in Pinot Noir grapevines. The difference observed in the 2006-2007 ABA application could possibly be due to a rain event that occurred a few hours after the ABA application, possibly diluting, washing off, or leaching out the applied ABA. We are repeating the ABA experiment this season to determine if ABA could be a possible tool in the management of Pierce's Disease.

**Status of Funds:** Approximately 1/3 of the total 2 year funding allocation was spent from 7/2007 to 2/2008. We anticipate spending the remaining 2/3 of the funds from 3/2008 to 6/2009.

**Summary and status of intellectual property produced during this research project:**

No intellectual property has been produced during this research period.