

Interim Progress Report for CDFA Agreement Number 12-0444-SA

Project Title: Field evaluation of grape plants expressing potential protective DNA sequences effective against Pierce's Disease.

Principal Investigator	David Gilchrist	Department of Plant Pathology, UC Davis	dggilchrist@ucdavis.edu
Co-Principal Investigator	James Lincoln	Department of Plant Pathology, UC Davis	jelincoln@ucdavis.edu
Collaborator	Mike Eldridge	Department of Plant Pathology, UC Davis	mdeldridge@ucdavis.edu
Collaborator	Abhaya Dandekar	Department of Plant Sciences, UC Davis	amdandekar@ucdavis.edu
Collaborator	Ann Powell	Department of Plant Sciences, UC Davis	ampowell@ucdavis.edu
Collaborator	Steven Lindow	Department of Plant and Microbial Biology, UC Berkeley	icelab@berkeley.edu

Reporting Period: These results are from work conducted from October 2015 to March 1, 2016

INTRODUCTION

The objective of this field experiment is to evaluate transgenic grape and grape rootstocks expressing various potentially protective genes from different constructs in a field site in Solano County for protection against Pierce's Disease (PD) caused by *Xylella fastidiosa* (Xf) (Pierce's Disease strain). Over the course of the multi-year field evaluation, test plants will include ungrafted conventional Thompson Seedless and Freedom plants as controls, transgenic plants from the Dandekar, Powell, Lindow and Gilchrist projects will evaluate the effect of several potential PD suppressive transgenes to protect the grape plants against symptom appearance and movement of the bacteria in the inoculated tissues. This experiment involving transgenic plants is permitted by APHIS and is regularly monitored for compliance to the permit specifications. The field experiment involves mechanical injections of Xf into the plant stems to establish infections. It is important to note that *X. fastidiosa* has existed in the same field area for the past 3 decades with no evidence of mechanical or vector spread from infected to adjacent healthy plants. Land preparation, planting, and management of the experimental resources to accommodate 500 plants was begun in 2010. The plants are on a row spacing of 15 feet between rows and 4 feet between plants in a row. The plants are irrigated by surface furrow in accordance with standard practices for maintaining grapes for experimental purposes at this site.

Although a drip irrigation system was installed in 2014 and will be used in all future plantings, furrow irrigation will be continued on the existing plots to avoid confounding of different water application methods in this experiment. Irrigation and pest management, primarily powdery mildew, weeds and insects, is coordinated by PI Gilchrist and conducted by Mike Eldridge the Field Superintendent employed by the Department of Plant Pathology who has 20 years' experience working locally with grapes and other perennial crops. The field crew work closely with PI Gilchrist to determine timing and need of each of the management practices.

OBJECTIVES

- A. Principal Investigators, with assistance from contract field crews, are responsible for pruning in the spring of each year and within the season as needed to maintain a reasonable canopy permitting sun exposure to leaves on inoculated canes. Periodic trimming is necessary, given that the transgenic plants are derived from Freedom (a common rootstock) and Thompson Seedless both of which exhibit tremendous vegetative growth during the season. In addition, annual pruning deviates from conventional practice in that multiple cordons have been established with a separate new cordon retained from each successive inoculation. This enables differential experimental materials for evaluation and sampling in the form of seasonal canes associated each succeeding annual inoculation. The objective is to provide sufficient inoculated and control material for destructive sampling over years to assess both timing of symptom development after successive inoculations and to assess bacterial presence and movement over time.
- B. Plants have been inoculated annually since 2011. *Xylella fastidiosa* was mechanically inoculated again in 2015. Sampling of inoculated stems from 2015 and previous years confirmed the presence of the bacteria and symptoms of Pierce's Disease later in the season.

DESCRIPTION OF ACTIVITIES TO ACCOMPLISH OBJECTIVES

The objectives set out for the establishment and management of this field planting were completed and the first planting occurred July 12, 2010 (Figure 1). The second phase of the planting, including grafted transgenics was completed May, 2011 and June of 2012.

Extensive polish trimming during the season is necessary to manage the Freedom and Thompson Seedless plants in a fashion to allow ease of mechanical inoculation and recovery of experimental samples (Figures 2, 3).

From planting through 2015 growing season, the individual transgenic plants all exhibited a normal phenotype, true to the untransformed control plants of each parental genotype (Figure 4). Symptoms of Pierce's Disease did not appear until two years after the first inoculation. Evaluations in the summer of 2014 and 2015 indicate all inoculated controls and some transgenic plants show symptoms of PD (Figure 5). It is clear that this field planting has provided important data on the effectiveness of any of the transgenic strategies employed by the respective researchers.

As of one year ago as the new buds began to emerge in mid to late March 2015, many inoculated canes on control plants and some transgenics did not survive the winter but the non-inoculated canes on these plants still appear healthy. Visual observation and destructive sampling of inoculated canes indicates that mechanical inoculation was successful in infecting inoculated canes (Figure 5). As of July 2015, in several uninoculated canes, adjacent to inoculated canes show foliar

symptoms indicting that the bacteria have moved systemically while some control plants are dead.

There are two points to be made regarding the appearance of symptoms. First, plant turgor has been maintained throughout the growing season with timely irrigation and there has been no evidence of wilt or epinasty symptoms prior to appearance of classic foliar symptoms or even immediately prior to death of inoculated control susceptible canes. Symptomatic leaves occur on inoculated canes without the appearance of water stress (Figure 5). This belies the long held anecdotal effect of vascular plugging leading to the classic foliar symptoms of sectorial death within green areas of leaves. Second, excellent symptoms associated with the presence of the pathogenic bacteria are readily seen in the spring of each year from buds emerging on inoculated canes. Buds break, push tiny leaves, and then die in tissues confirmed in the laboratory to harbor bacteria from inoculations that occurred one to two years prior.

There are now substantial differences between inoculated control plants compared with plants expressing some of the transgenes. There is no evidence of any spread of the bacteria from inoculated to non-inoculated control plants but there is now evidence of systemic spread within some of the plants representing different genetic composition (different transgenes). The positive result of effective mechanical inoculation over time suggests that plants consisting of transgenic root stocks grafted to non-transgenic scions will enable experimental assessment of cross-graft protection. Field data over the course of this experiment has been collected by all investigators and can be found in their individual reports from the 2014 and 2015 Pierce's Disease Symposium.

As of March 15th, 2015, all plants have been pruned to remove excess growth from the past year but to retain all inoculated wood (Figure 3). Spurs on old inoculated cordons were pruned to 2-3 buds while the 2014 inoculated branches were trimmed to retain up to 10 buds for data collection to include live/dead bud counting and destructive sampling for bacterial counts. The final inoculation was done in May 2015 and symptoms were observed on leaves from canes inoculated in 2014 (Figure 5).

We are approved and funded to continue maintenance and data collection from this site for the coming through June 30, 2016. This time period matches the time extension proposed by Dr. Dandekar, who has now assumed responsibility for the APHIS permit. Dr. Gilchrist will continue to manage the field operations at this site.

Field Work March to August 2015: All field activities are conducted or coordinated by field superintendent Mike Eldridge. Regular tilling and hand weeding maintained a weed-free planting area. Plants were pruned carefully in March leaving all inoculated/tagged branches and numerous additional branches for inoculation and sampling purposes in the coming year. All pruning material was left between the rows to dry, then flail chopped and later rototilled to incorporate the residue per requirements of the APHIS permit. Frequent trimming of the plants was done to ensure that leaves on inoculated canes were exposed to sunlight and shading of the associated leaves was avoided. Surface irrigation was applied as needed to maintain the soil at field capacity and turgor in the plants. Application of the fungicides Luna Experience and Inspire were alternated at periodic intervals to maintain the plants free of powdery mildew. Leafhoppers and mites were treated with insecticides when needed. Neither powdery mildew nor insect pressure was noted throughout the growing season. The same maintenance program is following the same general format for 2015 with timely pruning, rototilling weeds and applying fungicides on a monthly basis

to protect the vines against powdery mildew. Irrigation was applied at monthly intervals beginning in June. Pierce's disease symptoms were visible at bud emergence and later with foliar symptoms in late June and continuing to develop into late August.

Field Work from August 2015 to March 2016: the field was rototilled in the late fall after leaves had fallen and all Thompson Seedless grape clusters had been dropped.

RESEARCH RELEVANCE

The objective is to evaluate transgenic grape and grape rootstocks expressing various genes from different constructs in a field site in Solano County for protection against *Xylella fastidiosa* (Pierce's Disease strain) following mechanical injections of *X. fastidiosa* into the grape canes of both transgenic and co-planted non-transgenic control plants. Over the course of the multi-year field evaluation, test plants will include ungrafted conventional Thompson Seedless and Freedom plants as controls, transgenic plants from Dandekar, Powell, Lindow and Gilchrist projects will evaluate the effect of several potential PD suppressive transgenes to protect the grape plants against symptom appearance and movement of the bacteria in the inoculated tissues. The goal is to determine if these transgenes which suppressed PD symptoms in greenhouse experiments will also suppress in the field. All plants are located in an APHIS-approved field area with no risk of pollen or seed dispersal. The area is adjacent to experimental grape plantings that have been infected with Pierce's Disease for the past three decades with no evidence of spread of the bacteria to uninfected susceptible grape plantings within the same field plot.. The field area chosen has never had grapes planted therein, which is to avoid any potential confounding by soil borne diseases, including nematodes. Over the current 5 year course of this field experiment there is no evidence of spread of the bacteria to uninfected susceptible grape plantings.

LAYPERSONS SUMMARY

The purpose of the field planting is to evaluate grape and grape rootstocks expressing several transgenes from several investigators, with differing putative modes of action against *Xylella fastidiosa*, under natural field conditions for efficiency in providing protection against Pierce's Disease. The site in Solano County was selected and approved by APHIS to enable controlled inoculation and close monitoring of the host response in terms of symptoms, bacterial behavior, and plant morphology. Over the course of the multi-year field evaluation, test plants included ungrafted conventional Thompson Seedless and Freedom plants as controls, transgenic plants from investigators Dandekar, Labavitch, Lindow and Gilchrist and later transgenic rootstocks expressing some of the test genes were grafted to untransformed PD susceptible scions to assess potential for disease suppression in an untransformed scion from signals originating in the transformed rootstocks. We will continue maintaining and collecting data from this site for the coming season through June 30, 2016. This time period matches the time extension proposed by Dr. Dandekar, who has now assumed responsibility for the APHIS permit. Dr. Gilchrist will continue to manage the field operations at this site. The APHIS permit specifies that the plants are to be removed, burned on site and the field monitored as fallow for an additional year. This latter step may be modified if there is additional planting at this site. Additional space has been set aside to enable doubling of the current work area with a modification of the current permit.

Funding Agencies and Status of Funds: Funding for this project is provided by the CDFA Pierce's Disease and Glassy-winged Sharpshooter Board and the Regents of the University of California.

Intellectual Property: Evidence for any and all transgenes that show protection against PD will be submitted as a record of invention to the respective Technology Transfer offices at UC Davis and UC Berkeley as first step in protecting patent rights.

CONCLUSIONS

The results to date of this field experiment indicate that the mechanical inoculations successfully introduced the bacteria into the plants with subsequent appearance of foliar symptoms and cane death. There are transgenes from each of the investigators that appear to be suppressing the symptoms of PD inoculated vines.

Images below (Figures 1-5) illustrate the status the field experiment from planting in 2010 to the summer of 2015. The caption to each figure indicates when the image was obtained and together they represent the both asymptomatic inoculated transgenic and symptomatic inoculated non transgenic control plants at the Solano County site.

Publications: 2014 Pierce's Disease Symposium and 2015 Summary of Accomplishments posted on the CDFA Pierce's Disease website in lieu of a symposium-based report.

Images below show status of these plantings over the course of the experiment from early summer 2010 through July 2015



Figure 1 July 2010 Solano planting



Figure 2 July 2012 Solano planting



Figure 3. March 2015 Solano plants. Colored tags indicate site and date of inoculations 2011-2014:



Figure 4. Solano at the time of inoculation of the field plots with *Xylella fastidiosa*



Figure 5. Symptoms of Pierce's Disease recorded July 21, 2015 on branches inoculated May 27, 2014.